

Brain Signal Analysis

Advances in Neuroelectric and Neuromagnetic Methods

edited by Todd C. Handy

The MIT Press
Cambridge, Massachusetts
London, England

© 2009 Massachusetts Institute of Technology

All rights reserved. No part of this book may be reproduced in any form by any electronic or mechanical means (including photocopying, recording, or information storage and retrieval) without permission in writing from the publisher.

MIT Press books may be purchased at special quantity discounts for business or sales promotional use. For information, please email special_sales@mitpress.mit.edu or write to Special Sales Department, The MIT Press, 55 Hayward Street, Cambridge, MA 02142.

This book was set in Stone Sans and Stone Serif by SNP Best-set Typesetter Ltd., Hong Kong. Printed and bound in the United States of America.

Library of Congress Cataloging-in-Publication Data

Brain signal analysis : advances in neuroelectric and neuromagnetic methods / edited by Todd C. Handy.

p. cm.

Includes bibliographical references and index.

ISBN 978-0-262-01308-6 (hardcover : alk. paper)

1. Electroencephalography. 2. Magnetoencephalography. 3. Cognitive neuroscience—Methodology. I. Handy, Todd C.

QP376.5.B735 2009

616.8'047547—dc22

2008044258

10 9 8 7 6 5 4 3 2 1

Index

- AAHC (atomized and agglomerate hierarchical clustering), 37–38
 - and Krzanowski–Lai criterion, 39
 - for VEPs, 40, 41
- Action potentials, synchronization of, 179–81
- Adaptation hypothesis, 154
- Adaptive beamforming, 124
- Adjacent response (ADJAR) overlap estimation procedure, 218
- Aliasing, 176
- Alzheimer’s disease, 172
 - and synchronization likelihood, 192–93
- Amplitude
 - via analytic signal and Hilbert transform, 186–87
 - and volume conduction problem, 194
 - via wavelet analysis, 184–86
- Analytic signal, 186–87, 188
- ANOVA, 228
 - assumption of normality in, 235
 - and clustering, 41, 42
 - error rates in (type I), 234
 - repeated-measures, 226, 227, 228
 - topographic (TANOVA), 31–32, 42
 - and translational research, 226
- Argand plane, 176, 177
- Artifact(s), 26–27
 - in EEG, 214–15
 - and beamforming, 89
 - and data recording, 211, 220
 - detection procedures for, 215–17, 219
 - and instructions to participants, 212
 - in MEG, 106, 115–18
 - and memorization processes, 207
- Assumptions of covariance, in translational research, 228–29
- Assumptions of normality, in translational research, 235–37
- Assumption of sphericity, 228–29
 - in translational research, 228–29
- Atomized and agglomerate hierarchical clustering. *See* AAHC
- Attention, and MEG response amplitude, 115
- Attention, visual
 - and falls by elderly, 225
 - and VESPA method, 12, 13
- Attention search network, 171
- Autism
 - and sampling for translational research, 231
 - and synchronization disturbance, 172
- Auto-adaptive averaging, 216–17, 218, 219–20, 220
- Autoregressive techniques, 196
- Average reference, 26–27
- Averaging, 205–206, 219
 - auto-adaptive, 216–17, 218, 219–20, 220
 - grand average, 220–21
 - group-averaging, 43–44, 130–131
 - and signal-to-noise ratio, 113–114, 205–206

- Balanced design, 242
- Bayesian inference, 152
- Bayesian methods or approach, 141–42, 161
- Bayesian model comparison (selection), 149, 152–53, 158, 165
- Bayesian priors, 142–43, 151, 162
- Bayes's rule, 149, 151
- Beamformer spatial filter, 80–82, 96, 122–24
 - active and control conditions for, 88–89
 - adaptive, 124
 - bilateral, 83
 - and cancellation of correlated source activities, 82–87
 - and choice of head model, 91–92
 - and correlated activity, 124–26
 - and EEG, ix, 80, 82, 83, 123, 196, 198
 - group-averaging for, 130–31
 - linear (vector), 81
 - and localization, 195, 196
 - and MEG, ix, 81, 83, 94, 123, 133
 - and artifact reduction, 116
 - and correlated activity, 126
 - differential and event-related, 128–30
 - and SAM, 196, 198
 - minimum-variance, 133–36
 - and noise, 86, 87
 - number of EEG channels for, 90–91
 - number of trials for, 89–91
 - and overlap of MEG and EEG, ix
 - and signal to noise ratio, 126–28
 - specification of three-dimensional grid for, 93–94
 - statistical analyses for, 94–96, 131–33
 - time-frequency parameters for, 93
- BESA (Brain Electrical Source Analysis), 183, 196, 198
- BESA beamformer algorithms, 198
- BioSemi system, 155, 211, 212, 219
- Blind source separation techniques, 195
- Blurring, of temporal information, 91–92
- Bootstrapping (resampling strategy), 239, 244
- Brain Electrical Source Analysis (BESA), 183, 196, 198
- Brain function
 - and connectivity between regions, 171
 - functional specialization, 181–83
 - improved understanding of, 55
 - and local field potentials estimation, 55–56
 - (*see also* Local field potentials estimation)
 - through single-trial events, 56
- BRAINSTORM, 183
- Causation
 - and synchronization, 198–99
 - and topographic differences, 25, 33
- Cell population, activated by VESPA vs. VEP, 9–12
- Central limit theorem, 235
- Central tendency, robust measures of, 240
- Chaos theory, and synchronization
 - likelihood, 192–93
- Cluster analysis, 33–34
 - hierarchical clustering, 34, 37–38
 - k-means clustering, 34, 37
 - and optimal number of template maps, 37, 38–39
 - spatial correlation-based fitting in, 41–42
 - temporal considerations in, 39
- Coates, Roger, 177
- Cognition
 - cooperation required in, 171
 - and falls by elderly, 225
- Cognitive electrophysiology, and MRI, ix
- Cohen, David, 99
- Columnar oscillations, 179
- Complex numbers, 176
- Controls, between-group (translational research), 231–33
- Controls, within-group (translational research), 231
- Correlated source activities, and
 - beamforming, 82–87, 124–26
- Cortex, parietal, 73
- Coupling, of oscillators, 174
- Covariance, 235
 - assumptions of, 228–29

- Cross-sectional studies, 232
- Current source density (CSD), 25, 48n.4
- Data recording procedures, 211–12
- Data reduction, in dynamic causal modeling, 147–48
- Data resolution matrix, 65
- Data transforms, in translational research, 233
- DCM. *See* Dynamic causal modeling
- Differential beamformer, 128–29
- Dipole localization, zero, 61
- Dipole localization algorithms, 43
- Dipole models, 101, 103
- Dipole positioning, 79–80
- Dipole source localization
 - and DCM, 164
 - and number of electrodes, 90
- Distributed source estimation algorithms, 43
- Dynamic causal modeling (DCM), 142–43, 164–65
 - data reduction in, 147–48
 - event-related input and event-related response-specific effects in, 146–47
 - inference and model comparison in, 151–53
 - and mismatch negativity (MMN), 153–60
 - model priors in, 149, 150
 - and neural mass model, 143–46, 164
 - observation (likelihood) model, 148–49
 - and spatial forward model, 147
 - symmetry constraints in, 149, 151, 160–63
- Dynamic imaging of coherent sources (DICS), 196
- Dynamic SPM, 122
- Dyslexia
 - and magnocellular pathway function, 11
 - and synchronization disturbance, 172
- ECD model. *See* Equivalent current dipole model
- EcoG (electrocorticogram), 194
- EEG (electroencephalography), 141–42
 - artifacts in, 214–15
 - and average reference, 26–27
 - and Bayesian model comparison, 165
 - and beamformer, ix, 80, 82, 83, 123, 196, 198 (*see also* Beamformer spatial filter)
 - dynamic causal modeling (DCM) for, 142–46, 164–65 (*see also* Dynamic causal modeling)
 - and ECD modeling, 116
 - electrode placement in, 214
 - and ERPs, 45, 205
 - and functional or effective connectivity, 198
 - and global dissimilarity, 30
 - vs. hemodynamic techniques, 79
 - and inverse problem, 55, 79
 - and *k*-means clustering, 34, 37
 - and MEG, ix
 - and MRI, ix
 - noisy data for, 87
 - and participant factors, 105
 - research risk in, 46
 - and spatial smearing, 91
 - and synchronization, 175, 178, 182–83, 184, 187, 192–94 (*see also* Synchronization of neural oscillations)
 - traditional research on, 182
 - and VESPA, 5, 6
 - and visuomotor reaction time task, 73
 - and volume conduction problem, 193, 194, 195
- EEGLAB, 183
- Effective connectivity, 198
- ELECTRA source model, 55–56, 62, 67, 74
- Electrical Geodesic, Inc., 66
- Electrocaps, 214
- Electrocorticogram (EcoG), 194
- Electrode placement, 214
- Electrode preparation, 212
- Electroencephalography. *See* EEG
- Epilepsy, and synchronization likelihood, 192–93
- Epsilon estimate, 230, 233–34
- Equal means testing, 245
 - and ERP researchers, 226
 - generalizability of, 228
 - least squares approach to, 226

- Equivalent current dipole (ECD) model, 101, 122
 and dynamic causal modeling, 164
 and MEG artifact reduction, 116
 in Montreal Neurology Institute space, 157
- ERB (event-related beamforming) method, 126, 128, 129, 130, 131–32, 133
- ERFs (event-related fields), 109, 110, 113, 118
 and beamforming, 126, 127, 128, 130
 filtering of (auditory), 112
 and number of stimuli, 114, 115
 and number of trials, 113, 128
 topography of (auditory), 100
 visual, 117
- ERPs (event-related potentials), 1, 21–22, 205
 and cluster analysis, 33–46
 and dynamic causal modeling, 146
 and EEG, 45, 205
 error rates in data from, 226
 and global dissimilarity, 30
 and homogeneity of variance, 228
 and local field potentials estimation, 56
 and MRI, ix
 number of neural sources contributing to, 80
 and repeated-measures ANOVA, 226
 research risk in, 46
 shortcomings of data from, 56
 signal-to-noise optimizing for, 205–21 (*see also* Signal-to-noise ratio, optimizing of)
 and single-trial analysis, 74, 75
 topographic analysis of, 21, 44–46
 and traditional EEG research, 182
 and translational investigations, 225 (*see also* Translational ERP research)
- ERP waveforms
 and global field power, 29–30
 and reference dependence, 22–25
- ERR (event-related response), 146
- Error terms, pooled, 235
- Error terms, unpooled, 234–35
- ERS (event-related synchronization), 118–20
- Estimation procedures. *See also* Local field potentials (LFP) estimation; Source estimation
 ADJAR (adjacent response overlap estimation procedure), 218
 in DCM, 151
 for synchronous oscillations, 195–96
 of VESPA method, 15–16
- Euler, Leonard, 177
- Event-related beamforming (ERB) images, 126, 128, 129, 130, 131–32, 133
- Event-related fields. *See* ERFs
- Event-related potentials. *See* ERPs
- Event-related response (ERR), 146
- Event-related SAM, 129 (*see also* SAM)
- Event-related synchronization (ERS), 118–20
- Evoked fields, 113. *See also* Event-related fields
- Expectancy, and MEG response amplitude, 115
- Experiments, participants in. *See* Participants in experiments
- Factor levels, in translational-research design, 230–31
- Fast Fourier transforms (FFT), 111, 175, 178, 184
- FIELDTRIP, 183
- Filtering
 and data analysis operations, 217, 218–19
 as distorting temporal boundaries, 187
 in MEG, 111–12
 online, 212–13
 spatial, 122–23 (*see also* Beamformer spatial filter)
- Finger-tapping task, 66–73
- Fisher-scoring scheme, for dynamic causal modeling, 151
- Fitting, spatial correlation-based, 41–42
- FOCUSS, 122
- Fourier analysis
 of spectral power or phase, 175–79, 183
 and wavelet, 184

- Fourier series, 176–78
- Frontal cortex, and mismatch negativity, 154
- Functional brain networks, localization of, 55
- Functional connectivity, 198
- Functional magnetic resonance imaging (fMRI), 43, 55, 61, 99
- and beamformer analyses, 94–96
 - and cognitive electrophysiology, ix
 - and EEG, 79
 - and functional or effective connectivity, 198
 - and participant factors, 105
 - and translational principles, 228
- Functional microstates, principle of, 33, 39
- Functional specialization, in brain, 181–83
- Geodesics Netamps system, 66
- GEV (global explained variance), 34, 35, 37
- GFP (global field power), 27, 27–30
- Ghost sources, 43, 65, 74
- Glaucoma, and parvocellular and magnocellular systems, 11
- Global dissimilarity (DISS), 27, 28, 30–33
- Global explained variance (GEV), 34, 35, 37
- Global field power (GFP), 27, 27–30
- Goodness-of-fit tests, 237
- Gradiometers, 104
- planar vs. axial, 105
- Grand average, for ERPs, 220–21
- Granger causality, 196, 198–99
- Greenhouse–Geisser estimate, 233–34, 235
- Head models, 43, 56–57, 64, 68, 91–92, 93, 120–21, 147
- of Montreal Neurological Institute, 66
 - SYSMAC (symmetric spherical head model with anatomical constraints), 63
- Hemodynamic signal intensity, and translational principles, 228
- Hemodynamic techniques for examining brain function
- and beamformer images, 95–96
 - and distributed source estimation algorithms, 43
 - vs. EEG, 79
- See also* Functional magnetic resonance imaging; PET
- Hierarchical clustering, 34, 37–38
- AAHC, 37–38, 39, 40, 41
- Hilbert transform, 186–87
- Human Brain Mapping Consortium, 66
- Huygens, Christiaan, 172
- Huynh–Feldt estimate, 233–34, 235
- Independent component analysis (ICA), 195, 215
- Induced (rhythmic) fields, 113
- Inferotemporal (IT) cortex, 10
- Influence matrix, 65
- Information measures, of synchronization, 190–91
- Interpretability, neurophysiologic, and waveform reference, 23
- Inverse problem, 55, 58, 79, 194–95
- Inverse solution methods, 26
- and localization of single sources, 61
- Irotational source model, 60–62
- Kendall's tau, 228
- k*-means clustering, 34, 37
- Kolmogorov–Smirnov test, 237
- Krzanowski–Lai (*KL*) criterion, 36, 39
- Kullback–Leibler divergence, 152
- Kurtosis, 236, 237
- Laplacian waveforms, 25, 48n.4
- LAURA (algorithm), 43
- LCMV (linearly constrained minimum-variance) beamformer, 134
- MUSIC as, 123
- Lead field matrix, 81
- Least squares approach, 235
- and translational research, 226
- Left- and right-hand movements
- in finger-tapping task, 66–73
 - and visuomotor reaction-time task, 73
- Lehman, Dietrich, 25, 33, 39

- Linearly constrained minimum-variance (LCMV) beamformer, 134
- MUSIC as, 123
- Linear source estimation, 123
- Linear time-invariant (LTI) systems, 2
- Local field potentials (LFP), synchronization of, 179–81
- Local field potentials (LFP) estimation, 55–56
 - practical issues of, 62–63
 - and regularization, 64–65
 - statistical analysis on single trials, 65–66, 72–75
 - in spectral domain, 70–72
 - in temporal domain, 66–70
 - theoretical aspects of, 57–62
- Localization, source. *See at* Source localization
- Local synchronization, 181–83
- LORETA, 122

- Magnetoencephalography. *See* MEG
- Magnetometers, 104
- Magnocellular cells, and VESPA, 10–11
- MANOVA, 228, 235
- Matlab (MathWorks) code, 165
- Matlab routines, 183
- Matlab toolbox, 196
- Mauchly' s test, 229–30
- MEG (magnetoencephalography), 99–100, 133, 141–42
 - and artifacts, 106
 - removal and reduction of, 115–18
 - averaging in, 113–15
 - and Bayesian model comparison, 165
 - and beamforming, ix, 81, 83, 93–94, 123, 133
 - and artifact reduction, 116
 - body position for, 106–107
 - and correlated activity, 126
 - differential and event-related, 128–30
 - and SAM, 196, 198
 - data inspection and epoching for, 110–11
 - dynamic causal modeling (DCM) for, 142–46, 164–65 (*see also* Dynamic causal modeling)
 - and EEG, ix
 - filtering in, 111–12
 - and functional or effective connectivity, 198
 - head modeling and anatomical co-registration in, 120–22
 - head movement in, 107–108
 - head position for, 107
 - and hemodynamic techniques, 79
 - and inverse problem, 55, 79
 - and invisible sources of EEG, 62
 - and MRI, ix
 - neural basis of signal in, 100–103
 - and participant factors, 105–106
 - recording devices for, 103–105
 - recordings of, 109
 - signal analysis for, 109–10
 - stimulation procedures in, 109
 - and synchronization, 175, 178, 184, 187, 192–94 (*see also* Synchronization of neural oscillations)
 - and time-frequency response, 118–20
 - and VESPA, 5, 6
 - vigilance and task compliance in, 108, 110–11, 114
 - and volume conduction problem, 193–94
- MEG source estimation, beamformer for, 80
- M-estimators, 240, 241–42
- Migraine sufferers, 232
- Minimum norm solutions, 196
- Minimum-variance beamforming, 133–36
- Mismatch negativity (MMN) response, 114, 207
 - in dynamic causal modeling, 153–60
- Model-adjustment hypothesis, 154
- Model “inversion,” 141–42, 148, 166n.1
- Monitoring, online, 213–15
- Monte Carlo–Markov chain (MCMC) techniques, 165
- Montreal Neurological Institute
 - and group-average source estimations, 44
 - and locations of equivalent current dipoles, 157

- realistic head model from, 66
- template brain of, 131
- Morlet wavelet, 184
- MRI (magnetic resonance imaging), 106
 - for MEG head modeling, 121
- Multiple comparisons, in translation research, 243
- Multiple repetitions, and single-trial analysis, 56, 74
- Multiple simultaneous stimuli, and VESPA method, 12
- Multi-resolution analysis, 184–85
- Multisample sphericity, 229
- Multisensory integration processes, VESPA method in investigation of, 14
- MUSIC (beamformer), 123
- Mutual information, 191

- Neural activity index, 128, 135
- Neural information transmission, 100–103
- Neural mass model, 143–46, 164
- Neural synchronization, 172, 179, 180, 195–96. *See also* Synchronization of neural oscillations
- Neurocognitive disorder, translational investigation as help for, 225
- Neuron, as relaxation oscillator, 175
- Neurophysiologic interpretability, and waveform reference, 23
- Neuroprosthetics, 56
- NIRS (near-infrared spectroscopy), 99
- Noise. *See also* Signal-to-noise ratio, optimizing of
 - and beamformer spatial filter, 86, 87
 - phase slips from, 193
- Nonlinear modeling, 2, 3
- Normality
 - assumptions of (translational research), 235–37
 - tests of, 237
 - vs. variance, 227
- Novel sounds (“oddballs”), 153, 154, 155, 207

- Observation (likelihood) model (dynamic causal modeling), 148–49
- “Odballs” (novel sounds), 153, 154, 155, 207
- Online filtering, 212–13
- Online monitoring, during recording, 213–15
- Optimum allocation, 231
- Oscillations, columnar, 179
- Oscillators, 172–73
 - relaxation, 174–75
- Oscillatory synchronization, 174
- Outliers, 228, 236, 249, 242, 244–45

- Parietal cortex, 10, 73
- Participant factors, in MEG, 105–106
- Participants in experiments (subjects) and attributes of experiments, 210
 - vs. clinical population, 225–26
 - conduct toward, 211–12
 - reminder about instructions, 214
 - discarding (rejection) of, 220–21, 243
 - as drop-outs, 242
 - head models of, 91–92, 93, 120–21 (*see also* Head models)
 - heterogeneity (variability) among, 94, 225, 229
 - and MEG, 105–106
 - and artifacts, 106, 118
 - body position of, 106–107
 - head models of, 120–21
 - head position and movement of, 107–108, 115
- Parvocellular system, and VESPA, 10–11
- Performance data, and translational principles, 228
- Permutation tests, 237–39, 244
- PET (positron emission tomography), 43, 61, 99
 - and beamformer images, 96
 - and EEG, 79
 - and functional or effective connectivity, 198
 - and translational principles, 228

- Phase
 via analytic signal and Hilbert transform, 186–87
 Fourier analysis of, 175–79
 via wavelet analysis, 184–86
- Phase diffusion, 174
- Phase locking, 174
 gamma-band, 197
 in 1:1 vs. n:m ratio, 190–92
- Phase-locking value (PLV), 188
 normalized, 196
 and spurious synchronization, 193–95, 195–96
 and synchronization between neural sources, 195–96, 198
- Phase scattering, 189
- Phase slips, 174
- Phase synchronization, 172
- Physiological systems
 analysis of using stochastic stimuli, 1–2
 nonlinearities in, 3
- PLV. *See* Phase-locking value
- Pooled error terms, 235
- Positron emission tomography. *See* PET
- Prefrontal cortex, deviance detection system
 in, 154
- Principle component analysis, 116–18
- Proportionate allocation, 231
- Prospective studies, 232–33
- Pseudo-Z statistic, 128–29, 130, 135
- Quadratic analysis, and VESPA method, 8–9, 16
- Quasi-static approach, 57
- Quasi-stationarity, biophysical assumption of, 26
- Randomized control trials (RCTs), 231–32
- Receiver-operating characteristics (ROC), 123
- Recordings, of MEG, 109
- Recording time, maximum, 210
- Reference-dependence, of voltage ERP waveforms, 22–25
- Reference-independent spatial information.
See Topographic analysis methods
- Regions of interest (ROIs), anatomical, 95
- Regularization, and local field potentials estimation, 64–65
- Relaxation oscillators, synchronization of, 174–75
- Resampling strategies, 237, 242, 244
 bootstrapping, 239, 244
 permutation tests, 237–39, 244
- Retinitis pigmentosa, and parvocellular and magnocellular systems, 11
- Reverse correlation technique, 1
 and VESPA method, 3
- Right-hand rule, 100, 101, 103–104
- Robust measures, strategies of, 240, 242
 M-estimates, 241–42
 trimmed means, 240
 Winsorized means, 240, 240–41
- SAM (synthetic aperture magnetometry), 81, 126, 127, 196, 198
- Sampling, stratified, 231
- Sampling distribution, 236
- Scaling methods, 32
- Scalp current density (SCD), 194
- Scalp topography
 and location of brain generators, 183
 and reference dependence of waveforms, 25
 for VEP vs. VESPA, 9
- Schizophrenia
 early visual sensory processing deficit in, 11
 and magnocellular pathway function, 11
 and synchronization disturbance, 172
 and VESPA method, 12, 14
- Sessions, optimum number of, 210
- Shannon entropy, 191
- Signal analysis, for MEG, 109–10
- Signal-to-noise ratio, optimizing of, 205, 221
 and averaging, 113–14, 205–206 (*see also* Averaging)
 and beamformer, 87, 126–28
 collapsing across hemispheres, 208–209

- combining trial types, 206–207
- and data analysis
 - artifact detection, 215–17, 219
 - and order of operations, 217–20
- and data-recording procedures, 211–12
- and electrode preparation, 212
- by filtering, 111
 - online, 212–13
- grand average computed, 220–21
- and maximum recording time or number of sessions, 210
- number of conditions and stimulus positions optimized, 209
- and online monitoring during recording, 213–15
- proportion of stimuli of interest optimized, 207–208
- and signal analysis, 110
- trial length optimized, 209–10
- and variance, 244
- and VESPA vs. VEP, 5, 7
- Significance thresholds, of beamformer images, 132–33
- Singer, Wolf, 181
- Single-trial analysis, 44–46, 56
 - and local field potential estimation, 65–66, 72–75
 - in spectral domain, 70–72
 - in temporal domain, 66–70
- sLORETA, 43, 122
- SMA (spinal muscular atrophy), 70, 73, 74, 75
- SNR. *See* Signal-to-noise ratio, optimizing of
- Source estimation, 42–44, 48n.3. *See also*
 - Estimation procedures
 - and EEG channels, 219
 - linear techniques for, 122, 123
 - of MNI, 44
- Source localization, dipole
 - and DCM, 164
 - and number of electrodes, 90
- Source localization techniques or programs, 183–84
 - and oscillatory activation, 183
 - with synchronization and coherence analysis methods, 195–96
- Source modeling techniques, 80. *See also*
 - Beamformer spatial filter
- Spatial correlation (C), 27
- Spatial correlation-based fitting, 41–42
- Spatial domain, and single-trial analysis of
 - local field potentials, 67
- Spatial filter, beamforming as, 122–23. *See also* Beamformer spatial filter
- Spatial forward model, in dynamic causal modeling, 147
- Spatial smearing, 91, 107, 115
- Spatiotemporal analysis methods, 22
- Spatiotemporal beamformer, 129
- Spearman's rho, 228
- SPECT (single photon emission computed tomography), 61
- Spectral domain, and single-trial analysis of
 - local field potential, 70–72
- Spectral power, Fourier analysis of, 175–79
- Sphericity
 - assumption of, 228–29
 - tests of, 229–30
- SPM. *See* Statistical parametric mapping
- Spurious synchronization, 193–95
- Square inverse law, 61, 64
- Square-root rule of averaging, 113
- Statistical corrections, in translational research, 233–34
- Statistical nonparametric mapping (SnPM), 94
- Statistical parametric mapping (SPM), 94
 - dynamic SPM, 122
 - software for, 131, 152
- Statistical Parametric Mapping package (SPM8), 165
- Stochastic stimuli
 - in analysis of physiological systems, 1–2
 - in VESPA method, 3
- Strabismic amblyopia, and parvocellular and magnocellular systems, 11
- Stratified sampling procedures, 231

- Striate cortex, 10
 Superconducting quantum interference device (SQUID), 103–104
 Symmetric spherical head model with anatomical constraints (SYSMAC) method, 63
 Symmetry constraints, in dynamic causal modeling, 149, 151, 160–63
 Synchronization, 172
 and oscillation, 172–73, 174
 of relaxation oscillators, 174–75
 Synchronization of neural oscillations, 171–72
 of action potentials and local field potentials, 179–81
 complexity of interpretation of, 197
 local (and functional specialization), 181–83
 long-distance, 183–84
 phase and amplitude via analytic signal and Hilbert transform for, 186–87
 phase and amplitude via wave analysis for, 184–86
 measures of
 coherence, 192
 information measures, 190–91
 phase-locking value (PLV), 188–90
 spurious synchronization, 193–95
 synchronization likelihood, 192–93
 between neural sources, 195–98
 Synthetic aperture magnetometry (SAM), 81, 126, 127, 196, 198

 Talarach-Tourneoux, template brain of, 131
 TANOVA (topographic ANOVA), 31–32, 42
 Template maps, 34, 47
 optimal number of, 37, 38–39
 and spatial correlation-based fitting, 41–42
 temporal threshold for, 39
 in topographic pattern analyses, 41
 Temporal cortex, and mismatch negativity, 154
 Temporal domain, and single-trial analysis of local field potential, 66–70

 Temporal spectral evolution (TSE), 120
 Tests of normality, 237
 Tests of sphericity, 229–30
 Three-dimensional grid, for beamformer, 93–94
 Thresholding methods, for artifact detection, 215–16, 217, 219
 Time-frequency analysis, 93
 Time-frequency response (TFR) analysis, 118–20
 Topographic analysis methods, 21
 and average reference, 26–27
 and cluster analysis, 33–46
 for ERPs, 21, 44–46
 and global dissimilarity (DISS), 30–33
 and global explained variance, 35
 and global field power, 27–30
 and stable periods, 42
 for VEPs, 40–42
 Topographic ANOVA (TANOVA), 31–32, 42
 Topographic component recognition, 34
 Topographic differences, and causation, 25, 33
 Topographic pattern analyses. *See*
 Topographic analysis methods
 Topographic source estimations, 42–44
 Translational ERP research, 225–26
 analysis strategies for
 data transforms, 233
 statistical corrections, 233–34
 unpooled error terms, 234–35
 and assumptions of covariance, 228–29
 and assumptions of normality, 235–37
 and assumption of sphericity, 228–29
 challenges of, 244–45
 and covariance, 235
 design strategies for, 230
 controls (between-group), 231–33
 controls (within-group), 231
 on factor levels, 230–31
 multiple comparisons in, 243
 and resampling strategies, 237, 244
 bootstrapping, 239, 244
 permutation tests, 237–39, 244

- and robust-measures strategies, 240
 - M-estimates, 241–42
 - trimmed means, 240
 - Winsorized means, 240, 240–41
- and robust-measures strategies plus resampling, 242
- and tests of sphericity, 229–30
- and unbalanced designs, 242–43
- Trial length, optimizing of, 209–10
- Trimmed means, 240

- Unbalanced designs, 234, 242–43
- Unpooled error terms, 234–35

- Variance, 228
 - vs. normality, 227
 - and translational ERP studies, 244
- Vector scaling, 32
- Ventral visual stream, and VEP vs. VESPA, 10
- VEP (Visual evoked potential), 1, 22
 - and global field power, 30
 - topographic pattern analyses for, 40–42. 43
 - and VESPA, 3, 4, 5, 7, 8
 - and cell population (parvocellular and magnocellular), 9–12
- VESPA (visual evoked spread spectrum analysis) method, 1, 3–5, 14–15
 - acquisition of, 6
 - estimation of, 5, 15–16
 - extension of to higher orders, 8–9, 16
 - extension of to non-visual sensory systems, 12, 14
 - and multiple simultaneous stimuli, 12
 - and VEP, 3, 4, 5, 7, 8
 - and cell population (parvocellular and magnocellular), 9–12
- Visual attention. *See* Attention, visual
- Visual evoked potential. *See* VEP
- Volterra series, 2
- Volterra–Wiener approach, 2, 3
- Volume conduction problem, 193–94
- Voxel-based statistical results, 94

- Waveform analyses, 22
 - reference-dependence bias for, 22–25
- Wavelet analysis, 184–86
 - for phase-locking value, 188
- Weight sector normalized beamformers, 136
- Welch–James (*W*) statistic, 243
- White noise approach, 3
- Wiener, Norbert, 2
- Windowing, in MEG, 111
- Winsorized means, 240, 240–41

- z*-score, and PLV, 188
- z*-score transforms, 233